

Dynamical Mean Field Theory for strongly correlated electrons

TOOLbox series, nov 2021

Giacomo Mazza
DQMP, Unige

Monday 22

Introduction to the basics concepts of the Dynamical Mean Field Theory

(instruction for the hands-on tutorial)


Tuesday 23

Hands-on tutorial

Mott transition with iterated perturbation theory


Weak to strong coupling superconductivity
using Exact Diagonalization methods

General concepts of statistic physics
(single site representation, limit of infinite dimensions,..)




$$H_{\text{Hubbard}} = -t \sum_{\langle \mathbf{R}\mathbf{R}' \rangle \sigma} c_{\mathbf{R}\sigma}^\dagger c_{\mathbf{R}'\sigma} + U \sum_{\mathbf{R}} n_{\mathbf{R}\uparrow} n_{\mathbf{R}\downarrow} - \mu \sum_{\mathbf{R}\sigma} c_{\mathbf{R}\sigma}^\dagger c_{\mathbf{R}\sigma},$$

Lattice problem



Single-site representation



Compare with static mean-field

$$H_{\text{Ising}} = -J \sum_{\langle ij \rangle} \sigma_i \sigma_j - \sum_i h_i \sigma_i$$

	STATIC MeanField	DYNAMICAL MeanField
LOCAL OBSERVABLE	<p>local magnetization</p> $m_i = \langle \sigma_i \rangle$	
LOCAL REPRESENTATION	<p>spin in a magnetic field</p> $m_i = \tanh(\beta h_i^{\text{eff}})$	
SELF CONSISTENCY RELATION	$h_i^{\text{eff}} = h_i + J \sum_{\langle ij \rangle} m_j$	

	STATIC MeanField	DYNAMICAL MeanField
LOCAL OBSERVABLE	local magnetization $m_i = \langle \sigma_i \rangle$	local Greens function $G_{ii}(\tau) = -\langle T_\tau c_{i\sigma}(\tau) c_{i\sigma}^\dagger(0) \rangle$
LOCAL REPRESENTATION	spin in a magnetic field $m_i = \tanh(\beta h_i^{\text{eff}})$	atom in a bath $G_{ii}^{-1}(\tau) = \mathcal{G}_0^{-1}(\tau) - \Sigma_{\text{imp}}[\mathcal{G}_0]$
SELF CONSISTENCY RELATION	$h_i^{\text{eff}} = h_i + J \sum_{\langle ij \rangle} m_j$	$\Sigma_{\mathbf{k}}(\omega) = \Sigma_{\text{imp}}(\omega)$

$$\mathcal{S}_{\text{imp}}^{(i)} = \int d\tau d\tau' c_{i\sigma}^\dagger \mathcal{G}_0^{-1}(\tau - \tau') c_{i\sigma}(\tau') + \int d\tau H_{int}^{(i)}(\tau)$$

$$H_{int}^{(i)} = U n_{i\uparrow} n_{i\downarrow}$$

$$\mathcal{G}_0^{-1}(\tau - \tau') = G_{at,0}^{-1}(\tau - \tau') - \Delta(\tau - \tau').$$

$$G_{at,0}(i\omega_n) = i\omega_n + \mu$$

$$G_{ii,\sigma}(\tau - \tau') = -\langle T_\tau c_{i\sigma}(\tau) c_{i\sigma}^\dagger(\tau') \rangle_{\text{imp}[\mathcal{G}_0]}$$

$$G_{ii\sigma}^{-1}(\tau - \tau') = \mathcal{G}_0^{-1}(\tau - \tau') - \Sigma_{\text{imp}}[\mathcal{G}_0]$$

$$G_{ii}(\omega) = \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{\mathbf{k}}(\omega)}$$

DMFT approximation

$$\Sigma_{\mathbf{k}} = \Sigma_{\text{imp}}[\mathcal{G}_0]$$

exact for $z \rightarrow \infty$

initial guess for the Weiss field

$$\mathcal{G}_0$$

solve auxiliary impurity problem

get the impurity self-energy

get the local Greens function

get the new Weiss field

check for convergence

$$G_{imp}[\mathcal{G}_0]$$

$$G_{imp}^{-1}[\mathcal{G}_0] = \mathcal{G}_0^{-1} - \Sigma_{imp}[\mathcal{G}_0]$$

initial guess for the Weiss field

$$\mathcal{G}_0$$



solve auxiliary impurity problem



get the impurity self-energy



get the local Greens function



get the new Weiss field



check for convergence



$$G_{imp}[\mathcal{G}_0]$$

$$G_{imp}^{-1}[\mathcal{G}_0] = \mathcal{G}_0^{-1} - \Sigma_{imp}[\mathcal{G}_0]$$

$$\Sigma_{\mathbf{k}}(\omega) = \Sigma_{imp}(\omega)$$

$$G_{ii} = \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{imp}[\mathcal{G}_0]}$$

initial guess for the Weiss field

$$\mathcal{G}_0$$

solve auxiliary impurity problem

get the impurity self-energy

get the local Greens function

get the new Weiss field

check for convergence

$$G_{imp}[\mathcal{G}_0]$$

$$G_{imp}^{-1}[\mathcal{G}_0] = \mathcal{G}_0^{-1} - \Sigma_{imp}[\mathcal{G}_0]$$

$$\Sigma_{\mathbf{k}}(\omega) = \Sigma_{imp}(\omega)$$

$$G_{ii} = \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{imp}[\mathcal{G}_0]}$$

$$G_{ii} = G_{imp}$$
$$[\mathcal{G}_0^{\text{new}}]^{-1} = G_{ii}^{-1}[\mathcal{G}_0] + \Sigma_{imp}[\mathcal{G}_0]$$

initial guess for the Weiss field

$$\mathcal{G}_0$$



solve auxiliary impurity problem



get the impurity self-energy



get the local Greens function



get the new Weiss field



check for convergence



| $G_{imp}[\mathcal{G}_0]$

| $G_{imp}^{-1}[\mathcal{G}_0] = \mathcal{G}_0^{-1} - \Sigma_{imp}[\mathcal{G}_0]$

| $\Sigma_{\mathbf{k}}(\omega) = \Sigma_{imp}(\omega)$

| $G_{ii} = \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{imp}[\mathcal{G}_0]}$

| $G_{ii} = G_{imp}$
| $[\mathcal{G}_0^{new}]^{-1} = G_{ii}^{-1}[\mathcal{G}_0] + \Sigma_{imp}[\mathcal{G}_0]$

| $\mathcal{G}_0^{new} = \mathcal{G}_0$



initial guess for the Weiss field

$$\mathcal{G}_0$$

solver dependent!!

solve auxiliary impurity problem

get the impurity self-energy

get the local Greens function

get the new Weiss field

check for convergence

$$G_{imp}[\mathcal{G}_0]$$

$$G_{imp}^{-1}[\mathcal{G}_0] = \mathcal{G}_0^{-1} - \Sigma_{imp}[\mathcal{G}_0]$$

$$\Sigma_{\mathbf{k}}(\omega) = \Sigma_{imp}(\omega)$$

$$G_{ii} = \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{imp}[\mathcal{G}_0]}$$

$$G_{ii} = G_{imp}$$
$$[\mathcal{G}_0^{\text{new}}]^{-1} = G_{ii}^{-1}[\mathcal{G}_0] + \Sigma_{imp}[\mathcal{G}_0]$$

$$\mathcal{G}_0^{\text{new}} = \mathcal{G}_0$$

initial guess for the Weiss field

$$\mathcal{G}_0$$

solve auxiliary impurity problem

solver dependent!!

get the impurity self-energy

get the local Greens function

common to all
the solvers

get the new Weiss field

check for convergence

$$G_{imp}[\mathcal{G}_0]$$

$$G_{imp}^{-1}[\mathcal{G}_0] = \mathcal{G}_0^{-1} - \Sigma_{imp}[\mathcal{G}_0]$$

$$\Sigma_{\mathbf{k}}(\omega) = \Sigma_{imp}(\omega)$$

$$G_{ii} = \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{imp}[\mathcal{G}_0]}$$

$$G_{ii} = G_{imp}$$
$$[\mathcal{G}_0^{\text{new}}]^{-1} = G_{ii}^{-1}[\mathcal{G}_0] + \Sigma_{imp}[\mathcal{G}_0]$$

$$\mathcal{G}_0^{\text{new}} = \mathcal{G}_0$$

General Hamiltonian with local interactions

$$H = \sum_{\mathbf{R}\mathbf{R}'} \sum_{\alpha\beta} \sum_{\sigma} t_{\mathbf{R}\mathbf{R}'}^{\alpha\beta} c_{\mathbf{R}\alpha\sigma}^{\dagger} c_{\mathbf{R}'\beta\sigma} + \sum_{\mathbf{R}} H_{int}(\mathbf{R}).$$

$$\begin{aligned} H_{int}(\mathbf{R}) = & U \sum_{\alpha} n_{\mathbf{R}\alpha\uparrow} n_{\mathbf{R}\alpha\downarrow} + U' \sum_{\alpha \neq \beta} n_{\mathbf{R}\alpha\uparrow} n_{\mathbf{R}\beta\downarrow} + (U' - J_H) \sum_{\alpha < \beta} \sum_{\sigma} n_{\mathbf{R}\alpha\sigma} n_{\mathbf{R}\beta\sigma} \\ & + J_X \sum_{\alpha \neq \beta} c_{\alpha\uparrow}^{\dagger} c_{\beta\downarrow}^{\dagger} c_{\alpha\downarrow} c_{\beta\uparrow} + J_P \sum_{\alpha \neq \beta} c_{\alpha\uparrow}^{\dagger} c_{\alpha\downarrow}^{\dagger} c_{\beta\downarrow} c_{\beta\uparrow}. \end{aligned}$$

U intra-orbital repulsion

U' inter-orbital repulsion

J_H Hund's coupling

J_X Spin-flip

J_P Pair-hoppings

► Model Hamiltonians

► ab-initio Hamiltonian